


Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>DEC 2006</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Microbial Synthesis of 1,2,4-Butanetriol</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Chemistry Department, Michigan State University</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>Efforts were made to improve the microbial synthesis of D-1,2,4-butanetriol from D-xylose previously constructed (Scheme 1). Benzoylformate decarboxylase variants from Pseudomonas putida (ATCC 12633) with 2-fold increase in kcat/Km values were generated using directed evolution. Pseudomonas fragi D-xylonate dehydratase was purified and sequenced. E. coli yjhG and yagF encoding D-xylonate dehydratases, and yjhH and yagE genes encoding 3-deoxy-D-glycero-pentulosonate aldolases were identified, respectively. Site-specific knockouts of the aldolases prevented the cleavage of intermediate 3-deoxy-D-glycero-pentulosonic acid to form pyruvic acid and glycolaldehyde. D-Xylose dehydrogenase from Caulobacter crescentus CB15 was identified and expressed in E. coli that converts D-xylose into D-xylonic acid.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>SAR</b>	18. NUMBER OF PAGES <b>2</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

Final Technical Report  
Grant Number: N000140410134  
Principal Investigator: John W. Frost  
Performing Organization and Address: Chemistry Department, Michigan State University,  
E. Shaw Lane, East Lansing, MI 48824  
Grant Title: Microbial Synthesis of 1,2,4-Butanetriol  
Grant Period: January 1, 2003 – December 31, 2006

### Abstract/Project Summary:

Efforts were made to improve the microbial synthesis of D-1,2,4-butanetriol from D-xylose previously constructed (Scheme 1). Benzoylformate decarboxylase variants from *Pseudomonas putida* (ATCC 12633) with 2-fold increase in  $k_{cat}/K_m$  values were generated using directed evolution. *Pseudomonas fragi* D-xylonate dehydratase was purified and sequenced. *E. coli yjhG* and *yagF* encoding D-xylonate dehydratases, and *yjhH* and *yagE* genes encoding 3-deoxy-D-glycero-pentulosonate aldolases were identified, respectively. Site-specific knockouts of the aldolases prevented the cleavage of intermediate 3-deoxy-D-glycero-pentulosonic acid to form pyruvic acid and glycolaldehyde. D-Xylose dehydrogenase from *Caulobacter crescentus* CB15 was identified and expressed in *E. coli* that converts D-xylose into D-xylonic acid.

### Scheme 1.



### Scientific Technical Objectives:

Increasing key enzyme activity and eliminating side product biosynthesis will enhance the concentration and yield of D-1,2,4-butanetriol microbially synthesized from D-xylose. Directed evolution is widely used to generate enzyme variants with improved characteristics. Benzoylformate decarboxylase variants with substrate specificity towards 3-deoxy-D-glycero-pentulosonic acid will likely be identified. *E. coli* is known to use D-xylonic acid as a sole carbon source for growth, and the involved native pathway cleaves 3-deoxy-D-glycero-pentulosonic acid into pyruvate and glycolaldehyde. Identification and knockout of related genes becomes essential to increase D-1,2,4-butanetriol production. Microbial synthesis D-1,2,4-butanetriol from D-xylose requires D-xylose dehydrogenase activity for the conversion of D-xylose into D-xylonic acid. Identification of such enzyme enable the construction of a single *E. coli* microbe that is capable to convert D-xylose to D-1,2,4-butanetriol.

### Approach:

Error-prone PCR and DNA shuffling methods were used to create benzoylformate decarboxylase mutant library. A high throughput colorimetric assay was developed and used to screen for mutant with improved characteristics. D-Xylonate dehydratase from *P. fragi* was purified to homogeneity and its sequence was determined. This sequence in tandem with bioinformatics was to be used to identify D-xylonate dehydratase activity in *E. coli*. Based on the frequent clustering of genes in microbial catabolism, identification of this dehydratase activity led to the

identification of the gene encoding 3-deoxy-D-*glycero*-pentulose aldolase. Similarly, The partial DNA sequence of D-xylonate dehydratase was used to search the ERGO database to identify Orfs with 50 – 70% sequence homology. Identified Orfs in *Burkholderia fungorum* LB400 and *Caulobacter crescentus* CB15 were in close proximity to Orfs encoding a short chain dehydrogenase. These sequences were cloned, expressed in *E. coli* and assayed for D-xylose dehydrogenase activity.

### **Accomplishments:**

A high throughput screening method was developed to evaluate the *in vitro* benzoylformate decarboxylase variants towards 3-deoxy-D-*glycero*-pentulose. To generate the mutant library, the native *mdlC* gene encoding the wild-type benzoylformate decarboxylase from *P. putida* was first subjected to error-prone PCR. Mutant candidates with improved *in vitro* decarboxylase activities were then recombined using DNA shuffling. Mutant benzoylformate decarboxylases with up to 2-fold increase in  $k_{cat}/K_m$  values using 3-deoxy-D-*glycero*-pentulose as substrate were identified. Amino acid changes shared by these improved mutants were further identified by DNA sequencing. Genes *yjhG* and *yagF* were discovered to encode two isozymes of D-xylonate dehydratase in *E. coli*. Two isozymes of 3-deoxy-D-*glycero*-pentulose aldolase was also identified in *E. coli*, which are encoded by *yjhH* and *yagE*. Site-specific double knockout in the loci encoding the aldolase isozymes was generated. No growth was observed with this *E. coli* mutant when D-xylonic acid was used as the sole source of carbon. The genes encoding proteins RCO01012 from *C. crescentus* CB15 and RBU11704 from *B. fungorum* LB400 was discovered to possess D-xylose dehydrogenase activity. D-Xylose dehydrogenase from *C. crescentus* is the best candidate for expression in *E. coli* to create a construct capable of synthesizing D-1,2,4-butanetriol from D-xylose.

### **Significance:**

A practical route has been established enabling the synthesis of D-1,2,4-butanetriol from D-xylose.

### **Publications:**

None

### **Patent Information:**

Frost, J.W. U.S 2006/0234363, Oct. 19 2006.

Frost, J. W. WO 2008/091288, July 31, 2008.

### **Technology Transfer:**

Technology Licensed to Draths Corporation

### **Awards/Honors:**

None.